


Aug 17, 2022

Human Knee Meniscus Collection Protocol for scRNA-seq

DOI

dx.doi.org/10.17504/protocols.io.6qpvr614zvmk/v1



molmer¹, Martin Lotz¹

¹Scripps Research

Human BioMolecular Atlas Program (HuBMAP) Method Development Community
Tech. support email: Jeff.spraggins@vanderbilt.edu



Merissa Olmer
Scripps Research

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.6qpvr614zvmk/v1

Protocol Citation: molmer, Martin Lotz 2022. Human Knee Meniscus Collection Protocol for scRNA-seq. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.6qpvr614zvmk/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 15, 2022

Last Modified: August 17, 2022

Protocol Integer ID: 68682

Keywords: Meniscus, knee







Abstract

Meniscus is taken from the medial knee compartment is collected for scRNA-seq processing. The attached image indicates where the meniscal tissue is collected from.

Materials

Sterile drapes
Sterile gauze
Autoclaved tools
Disposable #21 scalpels
DPBS
DMEM
Antibiotic-Antimycotic
50mL conical tubes

Protocol materials

-  PAXgene Tissue STABILIZER **Qiagen Catalog #765512** Step 0.9
-  DMEM with L-Glutamine 4.5g/L Glucose and Sodium Pyruvate **Fisher Scientific Catalog #MT-10-013-CV** Step 4
-  Antibiotic-Antimycotic (100X) **Thermo Fisher Scientific Catalog #15240062** Step 4
-  PAXgene Tissue FIX **Qiagen Catalog #765312** Step 0.9

Before start


Knee blocks are shipped on wet ice and received within 48 hours post-mortem from certified tissue banks. The average time from receiving the sample to harvesting cartilage from both knees is about 2 hours. Pictures are taken of knee tissues and macroscopic scoring of the cartilage is completed.



- 1 Prepare harvesting area with sterile drapes, tools and gauze. All harvesting is completed within an aseptic environment.
- 2 Wipe down the knee blocks with 95% ethanol prior to opening the joint capsule.
- 3 Once the joint capsule is opened, the femur and tibia are disarticulated and menisci are resected.
- 4 A full-thickness (sagittal) piece of medial meniscus is cut out using a #21 scalpel; the piece measures about 1.60cmX15mmx8mm and weighs around 1g. This is collected into a 50mL tube that is filled with 40mLs of



DMEM with L-Glutamine 4.5g/L Glucose and Sodium Pyruvate **Fisher Scientific Catalog #MT-10-013-CV**

+ 1%  Antibiotic-Antimycotic (100X) **Thermo Fisher Scientific Catalog #15240062**, then placed in wet ice until the cell isolation protocol for scRNAseq begins.

Note

See Description Section for image for the area of interest.