

Brain pH Responses to Acetazolamide and Hypercapnia in Cats

Kiyotaka KOHSHI^{*,**}, Yoshimasa KINOSHITA^{*}, and Koichi FUKATA^{***}

Departments of ^{*}Neurosurgery and ^{**}Hyperbaric Medicine, School of Medicine,
and ^{***}Division of Clinical Examination, School of Medical Technology,
University of Occupational and Environmental Health, Kitakyushu, Fukuoka

Abstract

The involvement of increased brain tissue CO₂ tension in acetazolamide-induced brain acidosis was investigated by comparing the brain pH response to acetazolamide with that to hypercapnia. CO₂ and pH sensors were placed bilaterally into cerebral white matter to 15 mm depth in cats. Group 1 cats (n = 9) breathed spontaneously, and *in situ* brain tissue P_{CO₂} and pH (Pb_{CO₂} and pHb) were measured after intravenous acetazolamide administration (20 mg/kg). Group 2 cats (n = 9) were paralyzed and ventilated mechanically, and the changes of pHb were investigated by adjusting the ventilation to maintain the same Pb_{CO₂} values as in the acetazolamide-treated group. Pb_{CO₂} changes were not significantly different between the two groups. However, pHb responses were quite different: the fall in pHb was progressive in Group 1 but transient in Group 2. Brain acidosis after acetazolamide administration is not due to the rise in brain tissue CO₂ tension.

Key words: carbonic anhydrase, acetazolamide, pH regulation, cerebrovascular diseases

Introduction

Measurements of regional cerebral blood flow (CBF) have been used to evaluate the hemodynamic significance of ischemic cerebrovascular disease. Assessment of the capacity to increase CBF in response to vasodilators such as CO₂ or acetazolamide is another way of measuring cerebral circulatory reserve.

Acetazolamide, a selective inhibitor of carbonic anhydrase, is more easily administered to patients with cerebrovascular disease than CO₂, because it has little influence on the cardiopulmonary system.^{11,27} Acetazolamide increases the CBF remarkably and induces an increase in both brain tissue P_{CO₂} (Pb_{CO₂}) and carbonic acidosis, but the mechanisms for these effects have remained obscure. Moreover, there is a dissociation between the changes in Pb_{CO₂} and brain tissue pH (pHb); the fall in pHb is large despite the rise in Pb_{CO₂} being only slight after acetazolamide administration.^{4,28,30} The effects of acetazolamide on the acid-base balance might occur via a direct inhibition of carbonic anhydrase local-

ized on brain cells.^{4,30} However, the effects may be due to carbonic anhydrase inhibition in red blood cells (RBCs),^{21,22,31} because acetazolamide injected intravenously crosses the blood-brain barrier (BBB) slowly,^{21,23,29} whereas both Pb_{CO₂} and pHb change rapidly immediately after intravenous administration of acetazolamide.^{4,10,21,22,28} If these effects are caused only by disturbance of the CO₂-carrying capacity of RBCs, the pH responses to both acetazolamide and hypercapnia should be similar if the time course of Pb_{CO₂} is similar.

The recent development of the ion-sensitive field-effect transistor (ISFET) allows the measurement of P_{CO₂} and pH *in situ*.^{7,25} In animal and clinical experiments, catheter-tip CO₂ and pH sensors based on ISFET technology have been successfully used to monitor continuously *in situ* P_{CO₂} and pH in blood and tissues.^{18,25} The present study investigated whether acetazolamide-induced brain acidosis is due to the concomitant rise in brain tissue CO₂ tension.

Materials and Methods

Received April 17, 1996; Accepted November 26, 1996

Twenty-two cats of both sexes (body weight 3.4 ± 0.3 kg, mean \pm SD) were sedated with an in-

traperitoneal injection of 120 mg ketamine. Anesthesia was maintained by continuous infusion of ketamine (200 mg in 10 ml physiological saline) delivered at 2 ml/hr via a femoral venous catheter. The animals had miotic eyes and salivation during the measurements. Rectal temperature was monitored and kept at 38°C by means of a heating table. Arterial and venous catheters were inserted so that arterial P_{CO_2} (Pa_{CO_2}) could be monitored continuously and drugs could be infused. The animals were fixed to a stereotactic cradle, and a midline scalp incision was made after local anesthesia with 1% lidocaine. After the scalp was retracted, two holes of 0.5 cm diameter were drilled bilaterally through the cranium with a burr drill. Hemostasis was achieved with cauterization and bone wax. The centers of the holes were 1.0 cm anterior from the ears and 1.0 cm lateral from the midline. The exposed dura was coagulated and incised.

CO_2 and pH sensors (CO_2 sensor, CO-1035; pH sensor, pH-2135; Nihon Kohden, Tokyo) used catheter-tip electrodes which contained the pH-ISFET. The CO_2 sensor was coated with silicone, and the pH sensor was encapsulated in the end of a nylon tube. These CO_2 and pH sensors can measure continuously *in situ* P_{CO_2} and pH, respectively, in blood and other tissues.^{18,25)} The pH sensor reveals rapid responses to acidosis and alkalosis, and the response time of the CO_2 sensor is 2 minutes for a 90% change in CO_2 tension.²⁵⁾ The pH sensor was calibrated at 37°C with a standard phosphate buffer of pH 7.27, and the CO_2 sensor was calibrated at 37°C with sterile P_{CO_2} standard solutions of 36 and 86 mmHg. The baseline drift of the CO_2 sensor was less than 2 mmHg during 5 hours of measurements.¹⁸⁾ The CO_2 and pH sensors were inserted bilaterally to 15 mm depth from the cortical surface in the center of the holes. Another CO_2 sensor was inserted via a femoral arterial catheter to measure Pa_{CO_2} continuously. pH/ P_{CO_2} monitors (KR-5000; Nihon Kohden) were used to record the output of the CO_2 and pH sensors. Respiratory rate was monitored by the impedance method and averaged every 10 seconds, and heart rate was determined from the electrocardiogram.

Two groups of cats were studied. The first (Group 1) was designed to study the effects of acetazolamide and the second (Group 2) to study those of hypercapnia. Group 1 animals ($n = 9$) were allowed to breathe spontaneously for at least 30 minutes to let all parameters stabilize. At time 0 (control), after determination of all steady-state values, they received a bolus injection of 20 mg/kg acetazolamide dissolved in physiological saline (at a concentration of 100 mg/ml) via a femoral catheter. During the next 60

minutes, the animals remained in the cradle to monitor the parameters, and *in situ* Pb_{CO_2} and pHb were measured continuously. Group 2 animals ($n = 9$) underwent tracheotomy and muscle paralysis was achieved with intravenous pancuronium (1.0 mg/kg/hr). They were mechanically ventilated with air with a Harvard respirator (model 665d; Harvard Apparatus, South Natick, Mass., U.S.A.). Pb_{CO_2} was controlled by mechanical ventilation to maintain the mean Pa_{CO_2} value at that measured during the preacetazolamide period in Group 1 animals. Then, by reducing the rate and depth of respiration, Pa_{CO_2} was elevated to the mean Pb_{CO_2} value measured after acetazolamide injection within 10 minutes to reflect the Pb_{CO_2} change, because hypoxemia is not induced by hypercapnia^{20,33)} and the P_{CO_2} gradient between the brain and arterial blood is constant.¹⁶⁾ The mechanically induced Pa_{CO_2} elevation was maintained for more than 60 minutes. All parameters were recorded every 10 seconds by a programmable data logger (7V07; NEC San-Ei, Tokyo).

The obtained data were averaged over 1 minute (6 points) in each animal and analyzed using the Statview statistical package (Abacus Concepts). All values are reported as means \pm SE. Within each group, significant differences between time points were determined by one-way analysis of variance followed by Fisher's least significance test. For intergroup comparison of Pb_{CO_2} values, two-tailed unpaired *t* tests were applied. Statistical significance was taken as a *p* value of <0.05 .

Results

Satisfactory recordings of Pa_{CO_2} , Pb_{CO_2} , and pHb were obtained in 18 of 22 cats. The results in other animals, which developed bleeding at some point in the study, were discarded. The effects of acetazolamide and hypercapnia on Pa_{CO_2} , Pb_{CO_2} , and pHb are summarized in Table 1.

The changes in Pa_{CO_2} , Pb_{CO_2} , and pHb are illustrated by the representative electrode tracings from a Group 1 animal (Fig. 1). Acetazolamide administration was always followed by a rapid rise in Pb_{CO_2} , as well as by a concomitant fall in pHb. Such Pb_{CO_2} changes always reached a steady-state value by 20 minutes after drug administration, and always persisted for at least 40 minutes. The mean Pb_{CO_2} value increased by 25.3 ± 5.6 mmHg at 60 minutes after acetazolamide administration. In contrast, the fall in pHb gradually progressed during the measurements. pHb decreased by 0.20 ± 0.04 pH in 60 minutes. Pa_{CO_2} showed a transient slight rise immediately after acetazolamide administration and then decreased gradually, reaching a steady state at 10 minutes after

Table 1 Effects of acetazolamide and hypercapnia on $PbCO_2$ and pHb

Time (min)	Acetazolamide			Hypercapnia		
	$PbCO_2$ (mmHg)	pHb	$PaCO_2$ (mmHg)	$PbCO_2$ (mmHg)	pHb	$PaCO_2$ (mmHg)
-30	66.5 \pm 2.7	7.03 \pm 0.03	35.3 \pm 2.1	64.7 \pm 1.4	7.02 \pm 0.02	34.9 \pm 0.2
0	66.8 \pm 2.6	7.03 \pm 0.03	35.2 \pm 2.2	65.3 \pm 1.6	7.02 \pm 0.03	35.1 \pm 0.1
5	77.8 \pm 3.5	7.00 \pm 0.04	35.8 \pm 2.9	78.5 \pm 2.4	6.97 \pm 0.04	56.4 \pm 1.7
10	87.4 \pm 5.1	6.97 \pm 0.04	30.7 \pm 2.2	89.9 \pm 2.9	6.93 \pm 0.03	60.4 \pm 0.3
15	90.6 \pm 4.9	6.95 \pm 0.03	30.0 \pm 2.2	90.4 \pm 3.1	6.94 \pm 0.03	60.1 \pm 0.3
20	91.9 \pm 5.2	6.94 \pm 0.03	29.5 \pm 2.0	91.8 \pm 2.9	6.94 \pm 0.04	60.2 \pm 0.2
25	92.1 \pm 4.9	6.92 \pm 0.02	29.9 \pm 1.8	92.2 \pm 2.7	6.95 \pm 0.03	60.3 \pm 0.3
30	91.7 \pm 4.7	6.91 \pm 0.02	30.1 \pm 2.1	92.7 \pm 2.7	6.95 \pm 0.03	60.5 \pm 0.3
40	91.7 \pm 4.6	6.88 \pm 0.03	31.3 \pm 2.0	93.4 \pm 2.6	6.97 \pm 0.03	60.3 \pm 0.2
50	91.4 \pm 4.5	6.86 \pm 0.04	34.3 \pm 1.9	93.9 \pm 2.6	6.97 \pm 0.03	60.2 \pm 0.2
60	92.0 \pm 4.5	6.84 \pm 0.04	34.7 \pm 1.9	93.9 \pm 2.8	6.97 \pm 0.03	60.4 \pm 0.3
p	0.0001	0.0001	0.0001	0.009	0.0001	
Fisher	0/5-60	0/15-60	0/10-40	0/5-60	0/5-60, 10/40-60	

Values are means \pm SE. $PaCO_2$ and $PbCO_2$: arterial and brain tissue PCO_2 , pHb: brain tissue pH. Time 0 min indicates acetazolamide (20 mg/kg, i.v.) administration and reducing respiration. p and Fisher values were determined by analysis of variance.




Fig. 1 Representative $PaCO_2$, $PbCO_2$, and pHb changes in a Group 1 cat. Arrow indicates the time of intravenous administration of 20 mg/kg of acetazolamide.




Fig. 2 Representative $PaCO_2$, $PbCO_2$, and pHb changes in a Group 2 cat. Arrow indicates the time at which hypercapnia was induced.

drug infusion. The fall in $PaCO_2$ was maintained from 10 to 40 minutes. After this plateau phase, $PaCO_2$ increased gradually and reached a new steady-state value, which was not statistically different from the control value. No significant changes were found in respiratory and heart rates at any point.

Figure 2 shows representative tracings for $PaCO_2$, $PbCO_2$, and pHb in a Group 2 animal. The $PaCO_2$ was maintained at 35 ± 1 mmHg to match the $PbCO_2$ to the control $PbCO_2$ values in Group 1. $PaCO_2$ was increased by hyperventilation sufficient to increase $PbCO_2$ by 25 mmHg, i.e., the increase caused by

acetazolamide administration. pHb decreased from the baseline to the lowest value of 6.93 pH after 10-minute exposure to hypercapnia, and thereafter showed a significant rise from 40 minutes ($p < 0.05$). Heart rate decreased transiently after changing the ventilation and soon returned to the baseline.

The baseline values in $PbCO_2$ and pHb in the two groups were not significantly different. Reduction in the rate and depth of respiration changed $PbCO_2$ values similarly to those of the acetazolamide-treated animals. However, the time courses of pHb




Fig. 3 Mean percentage changes in $PbCO_2$ and pHb after intravenous 20 mg/kg acetazolamide administration.

changes in the two groups were quite different.

Discussion

This study found differences in both the trend and the magnitude of brain acidosis due to treatment with acetazolamide and hypercapnia despite maintenance of $PbCO_2$ at similar levels.

The time course of the mean percentage changes in $PbCO_2$ and pHb were quite different in animals treated with acetazolamide (Fig. 3), although the response time of the CO_2 sensor is slower than that of the pH sensor. This finding is in contrast to previous observations of the same mean percentage changes in P_{CO_2} and pH at the medullary surface.²⁸⁾ However, measurement of P_{CO_2} and pH at the brain surface found a fall in pH persisted whereas P_{CO_2} was reduced to the control values before treatment through changes in ventilation.⁴⁾ The reaction ($CO_2 + H_2O = H_2CO_3 = H^+ + HCO_3^-$) requires 20 seconds to reach 90% change of the total change in plasma.⁸⁾ The same time is estimated to be required to reach equilibrium in brain extracellular fluid because carbonic anhydrase is not present there. If both P_{CO_2} and pH are regulated together by this equilibrium reaction in brain extracellular fluid, then such a large difference in the mean percentage change would not occur even if the hydration reaction of CO_2 is uncatalyzed. Therefore, the previous and our present findings indicate that both brain H^+ and CO_2 can increase independently in response to acetazolamide.

The exact mechanisms by which acetazolamide induces a rise in $PbCO_2$ and brain acidification have not

been clarified. The importance of carbonic anhydrase in brain tissue has been emphasized in the rise in $PbCO_2$,⁴⁾ but carbonic anhydrase inhibition in RBCs causes both decreased CO_2 -carrying capacity of blood and greater P_{CO_2} gradient between tissue and blood,^{5,21,22,31)} because acetazolamide injected intravenously penetrates the BBB slowly and may take several hours before the physiological effect is maximum.^{21,23,29)} The overall pattern of carbonic anhydrase inhibition by acetazolamide injected intravenously is simply due to the greatly increased gradient of P_{CO_2} from tissue to alveolus.²¹⁾ Membrane-associated carbonic anhydrase is widely present on the luminal surface of endothelial cells of brain capillaries.⁹⁾ This type of carbonic anhydrase is present in the lungs³²⁾ and is involved in CO_2 exclusion from the lungs.⁶⁾ The carbonic anhydrase in brain capillaries is probably important for CO_2 transport from the brain to blood and inhibition may cause CO_2 retention in the brain.^{9,19)} Therefore, there are at least two sources for the rise in $PbCO_2$ after acetazolamide administration: inhibition of carbonic anhydrase located either in RBCs or in brain capillary endothelium.

Brain carbonic acidosis might occur via direct inhibition of carbonic anhydrase located in brain tissue, but the slow movement of acetazolamide across the BBB suggests that the rapid pHb change after drug administration is not caused by this mechanism. The excess rise in $PbCO_2$ after acetazolamide administration may cause the formation of carbonic acid in brain tissue.^{22,28)} According to this hypothesis, pHb responses to both acetazolamide and hypercapnia should be similar if the time course of $PbCO_2$ is the same. However, our study found the fall in pHb was progressive after acetazolamide administration, but transient in hypercapnia.

The pHb response to hypercapnia shows the phenomenon of pH buffering in brain extracellular fluid due to carbonic anhydrase in RBCs and brain tissue. Such brain pH buffering capacity is well known to be activated under hypercapnia and hypocapnia.^{1,2)} Our result that the pHb buffering is inhibited by acetazolamide administration suggests that one of the pHb regulating mechanisms is carbonic anhydrase-dependent. Brain pH is regulated mainly by transmembrane ion transport in both the brain capillary wall and the brain cell membrane.^{14,24)} Especially, Na^+/H^+ and Cl^-/HCO_3^- exchanges are important in brain pH regulation.^{3,12,14)} In the cerebrospinal fluid, Na^+/H^+ and Cl^-/HCO_3^- exchanges may be dependent on carbonic anhydrase located in the cells of the choroid plexus and glia.^{15,26)} Our result that acetazolamide inhibited pHb regulation is similar to the observation that in-

traperitoneal acetazolamide interfered with carbonic anhydrase-dependent ion transport in the brain cell membrane.¹⁷⁾ However, we believe that this phenomenon cannot be explained by this site of inhibition because of the slowness of acetazolamide penetration across the BBB. Intra- and extracellular pH changes are interdependent in brain tissue.¹³⁾ Thus, the inhibition of intracellular pH regulation might be greatly affected by the extracellular pH change. Although our data support the hypothesis that brain capillary endothelial carbonic anhydrase interferes with pH regulation in brain tissue, how intravenously injected acetazolamide affects the pH regulatory mechanism is not shown.

The pHb responses to acetazolamide and hypercapnia are quite different from each other when the values of P_{bCO_2} are not significantly different at any point during the measurement. Acetazolamide-induced brain carbonic acidosis is not due to the concomitant rise in brain tissue CO_2 tension.

References

- Adler S, Simplaceanu V, Ho C: Determination of rat brain buffering in vivo by ^{31}P -NMR. *J Appl Physiol* 64: 1829-1836, 1988
- Arief AI, Kerian A, Massry SG, DeLima J: Intracellular pH of brain: Alterations in acute respiratory acidosis and alkalosis. *Am J Physiol* 230: 804-812, 1976
- Bets AL: Sodium transport in capillaries isolated from rat brain. *J Neurochem* 41: 1150-1157, 1983
- Bickler PE, Litt L, Banville DL, Severinghaus JW: Effects of acetazolamide on cerebral acid-base balance. *J Appl Physiol* 65: 422-427, 1988
- Brzezinski J, Kjällquist Å, Siesjö BK: Mean carbon dioxide tension in the brain after carbonic anhydrase inhibition. *J Physiol (Lond)* 188: 13-23, 1967
- Effros RM, Chang RSY, Silverman P: Acceleration of plasma bicarbonate conversion to carbon dioxide by pulmonary carbonic anhydrase. *Science* 199: 427-429, 1978
- Esashi M, Matsuo T: Integrated micro multi ion sensor using field effect of semiconductor. *IEEE Trans Biomed Eng* 25: 184-192, 1978
- Forster RE, Crandall ED: Time course of exchanges between red cells and extracellular fluid during CO_2 uptake. *J Appl Physiol* 38: 710-718, 1975
- Ghandour MS, Langley OK, Zhu XL, Waheed A, Sly WS: Carbonic anhydrase IV on brain capillary endothelial cells: A marker associated with the blood-brain barrier. *Proc Natl Acad Sci U S A* 89: 6823-6827, 1992
- Heuser D, Astrup J, Lassen NA, Betz E: Brain carbonic acid acidosis after acetazolamide. *Acta Physiol Scand* 93: 385-390, 1975
- Højer-Pedersen E: Effect of acetazolamide on cerebral blood flow in subacute and chronic cerebrovascular disease. *Stroke* 18: 887-891, 1987
- Javaheri S, De Hemptinne A, Vanheel B, Leusen I: Changes in brain ECF pH during metabolic acidosis and alkalosis: A microelectrode study. *J Appl Physiol* 55: 1849-1853, 1983
- Jean T, Frelin C, Vigne P, Lazdunski M: The Na^+/H^+ exchange system in glial cell lines: Properties and activation by an hyperosmotic shock. *Eur J Biochem* 160: 211-219, 1986
- Katsura K, Kristián T, Nair R, Siesjö BK: Regulation of intra- and extracellular pH in the rat brain in acute hypercapnia: A re-appraisal. *Brain Res* 651: 47-56, 1994
- Kazemi H, Choma L: H^+ transport from CNS in hypercapnia and regulation of CSF $[HCO_3^-]$. *J Appl Physiol* 42: 667-672, 1977
- Kennealy JA, McLennan JE, Loudon RG, McLaurin RL: Hyperventilation-induced cerebral hypoxia. *Am Rev Respir Dis* 122: 407-412, 1980
- Kjällquist Å, Messeter K, Siesjö BK: The in vivo CO_2 buffer capacity of rat brain tissue under carbonic anhydrase inhibition. *Acta Physiol Scand* 78: 94-102, 1970
- Kohama A, Nakamura Y, Nakamura M, Yano M, Shibata K: Continuous monitoring of arterial and tissue P_{CO_2} . *Crit Care Med* 12: 940-942, 1984
- Kohshi K, Konda N, Kinoshita Y, Tsuru E, Yokota A: In situ arterial and brain tissue P_{CO_2} responses to acetazolamide in cats. *J Appl Physiol* 76: 2199-2203, 1994
- Kohshi K, Yokota A, Konda N, Kinoshita Y, Kajiwarra H: Intracranial pressure responses during hyperbaric oxygen therapy. *Neurol Med Chir (Tokyo)* 31: 575-581, 1991
- Maren TH: Carbonic anhydrase: Chemistry, physiology, and inhibition. *Physiol Rev* 47: 595-781, 1967
- Meyer JS, Gotoh F: Interaction of cerebral hemodynamics and metabolism. *Neurology* 11: 46-65, 1961
- Roth LJ, Schooler JC, Barlow CF: Sulfur- 35 labeled acetazolamide in cat brain. *J Pharmacol Exp Ther* 125: 128-136, 1959
- Seifter JL, Aronson PS: Properties and physiologic roles of the plasma membrane sodium-hydrogen exchanger. *J Clin Invest* 78: 859-864, 1986
- Shimada K, Yano M, Shibata K, Komoto Y, Esashi M, Matsuo T: Application of catheter-tip i.s.f.e.t. for continuous in vivo measurement. *Med Biol Eng Comput* 18: 741-745, 1980
- Siesjö BK, Kjällquist Å: A new theory for the regulation of the extracellular pH in the brain. *Scand J Clin Lab Invest* 24: 1-9, 1969
- Sullivan HG, Kingsbury TB IV, Morgan ME, Jeffcoat RD, Allison JD, Goode JJ, McDonnell DE: The rCBF response to Diamox in normal subjects and cerebrovascular disease patients. *J Neurosurg* 67: 525-534, 1987
- Teppema LJ, Rochette F, Demedts M: Effects of acetazolamide on medullary extracellular pH and P_{CO_2} and on ventilation in peripherally chemodenerated cats. *Pflügers Arch* 415: 519-525, 1990

- 29) Travis DM, Wiley C, Maren TH: Respiration during chronic inhibition of renal carbonic anhydrase: Further observations on pharmacology of 2-benzene-sulfonamido-1,3,4-thiadiazole-5-sulfonamide (CL 11366), acetazolamide and methazolamide. *J Pharmacol Exp Ther* 151: 464-481, 1966
- 30) Vorstrup S, Brun B, Lassen NA: Evaluation of the cerebral vasodilatory capacity by the acetazolamide test before EC-IC bypass surgery in patients with occlusion of the internal carotid artery. *Stroke* 17: 1291-1298, 1986
- 31) Vorstrup S, Henriksen L, Paulson OB: Effect of acetazolamide on cerebral blood flow and cerebral metabolic rate for oxygen. *J Clin Invest* 74: 1634-1639, 1984
- 32) Waheed A, Zhu XL, Sly WS: Membrane-associated carbonic anhydrase from rat lung. *J Biol Chem* 267: 3308-3311, 1992
- 33) Zauner A, Bullock R, Di X, Young HF: Brain oxygen, CO₂, pH, and temperature monitoring: Evaluation in the feline brain. *Neurosurgery* 37: 1168-1177, 1995

Address reprint requests to: K. Kohshi, M.D., Departments of Neurosurgery and Hyperbaric Medicine, School of Medicine, University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu, Fukuoka 807, Japan.

Commentary

Kohshi et al. have provided important insight into the mechanism of induction of brain acidosis by acetazolamide. In their well designed and creative set of experiments, these investigators have shown clear differences in the temporal profile and magnitude of brain acidosis due to acetazolamide and hypercapnia. By adjusting for Pb_{CO_2} in the two experimental groups, they have shown that the mechanism for acetazolamide-induced brain acidosis is not simply the result of a rise in brain CO₂ tension induced by the pharmacologic agent.

As with most good experiments, this study answers an important question and also raises some new unanswered questions that require further investigation. The authors' scholarly discussion applies their novel finding to existing hypotheses on mechanisms by which acetazolamide induces a rise in Pb_{CO_2} and brain acidification. The results strongly suggest that brain capillary endothelial carbonic anhydrase interferes with pH regulation in brain tissue. Further work is indicated to determine the precise mechanism by which acetazolamide affects the pH regulatory mechanism.

Daniel L. BARROW, M.D.
Department of Neurosurgery
The Emory Clinic

Georgia, U.S.A.

Dr. Kohshi and his colleagues report interesting data which disclose striking differences in brain tissue pH responses between acetazolamide administration and hypercapnia. They demonstrate that, when acetazolamide is administered intravenously, the brain tissue pH decreases slowly, whereas the increase in CO₂ tension of the brain tissue reaches its maximum level far more rapidly. In contrast, hypercapnia causes a decrease in pH in parallel with an increase in CO₂ tension within the brain tissue. This difference in immediate responses of the brain tissue pH should be taken into account when cerebrovascular responses are analyzed by these procedures. They also demonstrate that, while the decrease in brain tissue pH during hypercapnia is transient, acetazolamide causes the brain tissue pH to decrease progressively. As mentioned by the authors, the slowly progressive decrease in brain tissue pH caused by acetazolamide may reflect disturbances in the mechanisms of pH regulation. The authors suggest that inhibition of carbonic anhydrase located on the capillary endothelial cells of the brain could play a major role in producing such disturbances of pH regulation within the brain tissue. This is another important factor which should be borne in mind when acetazolamide is used clinically.

Yoichi KATAYAMA, M.D.
Department of Neurological Surgery
Nihon University School of Medicine
Tokyo, Japan

Kohshi et al. have used CO₂ and pH sensors, which facilitate continuous monitoring of brain tissue CO₂ tension (Pb_{CO_2}) and pH (pHb), and studied the effects of acetazolamide and hypercapnia on Pb_{CO_2} and pHb. It is well-known that acetazolamide is a potent cerebrovascular dilator and it is used to examine cerebrovascular reserve capacity in stroke patients. However, the mechanism by which acetazolamide causes tissue acidosis is not completely understood. The authors demonstrated clear Pa_{CO_2} , Pb_{CO_2} , and pHb changes in acetazolamide and hypercapnia groups. Acetazolamide-treated animals showed progressive and persistent falls in pHb associated with Pb_{CO_2} rise. Hypercapnia animals showed the same Pb_{CO_2} rise as seen in acetazolamide-treated animals, but the fall in pHb was transient. Thus, the pHb responses to acetazolamide and hypercapnia are quite different when the values of Pb_{CO_2} are not significantly different at any point during the measurement. They concluded that acetazolamide-induced brain carbonic acidosis was not due to the concomitant rise in brain tissue CO₂ tension. This conclu-

sion rises another question about the discrepancy between the rapidity of the brain acidosis after acetazolamide administration and the slowness of acetazolamide penetration across the BBB. Further new information about the biochemical effects of acetazolamide on cerebral pH regulation are necessa-

ry to answer these complex problems.

Minoru MAEDA, M.D.

Department of Neurosurgery
Juntendo University Izunagaoka Hospital
Shizuoka, Japan